

In the specification:

On page 12, please amend the paragraphs beginning at line 10 as follows:

A preferred, non-limiting example of such a mathematical algorithm is described in Karlin *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., NBLAST) can be used. See ~~http://www.ncbi.nlm.nih.gov~~. In one embodiment, parameters for sequence comparison can be set at score = 100, wordlength = 12, or can be varied (e.g., W = 5 or W = 20).

In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman *et al.* (1970) (*J. Mol. Biol.* 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package (~~available at~~ ~~http://www.gcg.com~~), using either a BLOSUM 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux *et al.* (1984) *Nucleic Acids Res.* 12(1):387) (~~available at~~ ~~http://www.gcg.com~~), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6.

On page 20, please amend the paragraph beginning at line 2 as follows:

The protein sequences of the present invention can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997)

*Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See ~~http://www.ncbi.nlm.nih.gov.~~

On page 39, please amend the paragraph beginning at line 14 as follows:

The nucleotide sequences of the present invention can be used as a "query sequence" to perform a search against public databases, for example, to identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See ~~http://www.ncbi.nlm.nih.gov.~~